

**TREATMENT OF PREECLAMPSIA AND PRETERM  
LABOR WITH COMBINATION OF PROGESTATIONAL AGENT  
AND A NITRIC OXIDE SYNTHASE SUBSTRATE AND/OR DONOR**

**Background of the Invention**

5 This invention relates to a method for the treatment of preeclampsia and of preterm labor with the combination of a progestational agent and a nitric oxide synthase substrate, a nitric oxide donor or both, alone or in further combination with one or more of a cyclooxygenase inhibitor, a PGI<sub>2</sub>-mimetic, a thromboxane (TXA<sub>2</sub>) inhibitor, 10 A compound possessing TXA<sub>2</sub>-agonistic and TXA<sub>2</sub>-inhibiting properties, a compound possessing TXA<sub>2</sub> antagonistic and PGI<sub>2</sub>-memetic activities, and a TXA<sub>2</sub> antagonist, and to pharmaceutical compositions comprising such a 15 combination.

Preeclampsia, toxemia or eclampsia of pregnancy can be a significant health problem during pregnancy and they are the leading causes of fetal growth retardation, fetal mortality and morbidity, premature birth and maternal 20 mortality. The etiology of the disease is largely unknown and effective therapy is not available. Preeclampsia of pregnancy is characterized by a triad of hypertension, pathological edema and proteinuria. This disease affects 6 to 10% of all pregnancies.

25 Recently, nitric oxide has been shown to be endothelium derived relaxing factor (EDRF) from the endothelium of blood vessels. Nitric oxide is considered to be a major mediator in the control of vascular reactivity. Nitric oxide is synthesized from L-arginine by nitric 30 oxide synthase located in endothelial cells. Nitric Oxide can also be generated by application of various nitric oxide donors such as sodium nitroprusside, nitroglycerin, glyceryl trinitrite, SIN-1, isosorbide mono-nitrite, isosorbide dinitrite, etc.

35 Treatment of pregnant rats with nitric oxide synthase inhibitors, which are analogues of L-arginine (such as L-NAME, N<sup>G</sup>-nitro-L-arginine methyl ester) results in

elevated blood pressure, fetal retarded growth and proteinuria. Thus, inhibition of nitric oxide synthesis produces conditions and symptoms identical to preeclampsia of pregnancy and establishes that preeclampsia is the direct result of the decrease in nitric oxide synthesis and/or a change in the regulation of vascular tone. These conditions give rise to increased blood pressure, decreased blood flow to the fetus, retarded fetal development and proteinuria. Agents which raise nitric oxide levels therefore are useful in the treatment of preeclampsia of pregnancy. Since nitric oxide donors also reduce contractility of the uterus during pregnancy, nitric oxide donors are also useful for use in preterm labor.

The nitric oxide effects on smooth muscle depend upon the activation of guanylate cyclase and generation of cGMP to produce relaxation and this step is progesterone dependent. Thus, combinations of nitric oxide donors with progesterone are particularly efficacious for the treatment of preeclampsia and of preterm labor.

EP 0 441 119 A2 discloses the use of L-arginine in the treatment of hypertension and other vascular disorders. It suggests that the mechanism by which L-arginine is effective for this purpose is because it may be the physiological precursor of "the most powerful endothelial-derived releasing factor, nitric oxide." The use of L-arginine in combination with other pharmaceutically active agents is not discussed in this publication.

#### Objects of the Invention

It is an object of the invention to provide a method for the prevention and treatment of preeclampsia with a combination of a progestational agent and a nitric oxide substrate and/or donor.

It is another object to provide such a method in which a progestational agent is used in combination with

a nitric oxide substrate and/or donor for the prevention and treatment of preeclampsia.

5 It is a further object to provide a method for the prevention and treatment of preterm labor using a pro-gestational agent in combination with a nitric oxide substrate and/or donor.

A further object is the provision of pharmaceutical compositions useful in practicing the methods of this invention.

10 Other objects will be apparent to those skilled in the art to which this invention pertains.

Summary of the Invention

15 In a method aspect, this invention relates to a method of treating at least one of preeclampsia and pre-term labor in a pregnant female which comprises administering to a pregnant female manifesting the symptoms thereof, (a) a progestational agent and (b) one or both of a nitric oxide synthase substrate and a nitric oxide donor, alone or in further combination with one or more  
20 of a cyclooxygenase inhibitor, a  $\text{PGI}_2$ -mimetic, a thromboxane ( $\text{TXA}_2$ ) inhibitor, a compound possessing  $\text{TXA}_2$ -agonistic and  $\text{TXA}_2$ -inhibiting properties, a compound possessing  $\text{TXA}_2$ -antagonistic and  $\text{PGI}_2$ -mimetic activities, and a  $\text{TXA}_2$  antagonist, in amounts effective to ameliorate the  
25 symptoms thereof, the amount of the progestational agent administered being bioequivalent to 50-300 mg. of injected progesterone and the amount of the nitric oxide synthase substrate, nitric oxide donor or both being effective to, respectively, either raise the blood level  
30 of circulating L-arginine in a pregnant female to whom the composition is administered to at least about 1 mmole above the normally 2 to 3 mmolar circulating levels or raise nitric oxide donor levels to about 1 to 100 nmolar (nanomolar).

35 In another method aspect, this invention relates to a method of treating preterm labor in a pregnant female which comprises administering to a pregnant female

manifesting the symptoms thereof, amounts of (a) a pro-  
gestational agent and (b) at least one of a nitric oxide  
synthase substrate and a nitric oxide donor effective to  
terminate the preterm labor, alone or in further combina-  
5 tion with one or more of a cyclooxygenase inhibitor, a  
PGI<sub>2</sub>-mimetic, a thromboxane (TXA<sub>2</sub>) inhibitor, a compound  
possessing TXA<sub>2</sub>-agonistic and TXA<sub>2</sub>-inhibiting properties,  
a compound possessing TXA<sub>2</sub>-antagonistic and PGI<sub>2</sub>-memetic  
activities, and a TXA<sub>2</sub> antagonist, the amount of the pro-  
10 gestational agent administered being bioequiva-lent to  
50-300 mg. of injected progesterone and the amount of the  
nitric oxide synthase substrate, nitric oxide donor or  
both being effective to, respectively, either raise the  
blood level of circulating L-arginine in a pregnant fe-  
15 male to whom the composition is administered to at least  
about 1 mmole above the normally 2 to 3 mmolar circulat-  
ing levels, or raise nitric oxide donor levels to about 1  
to 100 nmolar.

In a product aspect, this invention relates to a  
20 pharmaceutical composition comprising (a) a progesta-  
tional agent and (b) at least one of a nitric oxide  
synthase substrate and a nitric oxide donor, alone or in  
further combination with one or more of a cyclooxygenase  
inhibitor, a PGI<sub>2</sub>-mimetic, a thromboxane (TXA<sub>2</sub>) inhibitor,  
25 a compound possessing TXA<sub>2</sub>-agonistic and TXA<sub>2</sub>-inhibiting  
properties, a compound possessing TXA<sub>2</sub>-antagonistic and  
PGI<sub>2</sub>-memetic activities, and a TXA<sub>2</sub> antagonist, with the  
amount of the progestational agent per unit dosage being  
bioequivalent to 50-300 mg. of injected progesterone and  
30 the amount of the nitric oxide synthase substrate, a  
nitric oxide donor or both per unit dosage being  
effective to, repsectively, either raise the blood level  
of circulating L-arginine to at least about 1 mmole above  
the normally 2 to 3 mmolar circulating levels or raise  
35 the nitric oxide donor levels to about 1 to 1000 nmolar.

Detailed Disclosure

The methods of this invention treat one or more of preeclampsia and preterm labor in a pregnant female mammal, preferably a human, who is manifesting the symptoms thereof or who is a high risk candidate for doing so, e.g., as determined by the progress of a present or previous pregnancy.

Because these abnormal conditions of pregnancy are produced by or aggravated by subnormal nitric oxide synthesis, both nitric oxide synthase substrates, e.g., L-arginine, and nitric oxide donors, e.g., sodium nitroprusside, nitroglycerin, glyceryl trinitrate, SIN-1, isosobid mononitrate and isosorbid dinitrate, are useful for ameliorating the symptoms thereof and, in one aspect of the method of this invention, a combination of both are employed.

A synergistic effect is achieved when a progestational agent is administered concurrently with the nitric oxide substrate and/or nitric acid donor.

Thus, the method aspect of this invention and the pharmaceutical composition aspect of this invention employs a combination of (a) a progestational agent, e.g., progesterone, and (b) either or both of a nitric oxide donor and a nitric oxide synthase substrate and, optionally, (c) one or more of a cyclooxygenase inhibitor, e.g., aspirin; a PGI<sub>2</sub>-mimetic, e.g., iloprost and cica-prost; a thromboxane (TXA<sub>2</sub>) inhibitor, e.g., dazoxiben hydrochloride (benzoic acid, 4-[2-(1H-imadazol1-yl)-ethoxy]-, monohydrochloride; UK 37248), dazmegrel (1H-indole-1-propanoic acid, 3-(1H-imidazol-1-ylmethyl)-2-methyl-; UK 3885), ozagrel (2-propenoic acid, 3-[4-(1H-imidazol-1-ylmethyl)phenyl]-; OKY-046) and pirmagrel (imidazo[1,5-a]pyridine-5-hexanoic acid; CGS-13080); a compound possessing TXA<sub>2</sub>-agonistic and TXA<sub>2</sub>-inhibiting properties, e.g., ridogrel (pentanoic acid, 5-[[[3-pyridinyl[3-(trifluoromethyl)phenyl]methylene]-amino]oxy]-; R-68070) and labogrel (6-heptenoic acid, 7-phenyl-7-(3-

pydridinyl)-; a compound possessing TXA<sub>2</sub>-antagonistic and PGI<sub>2</sub>-memetic activities, e.g., 5-heptenoic acid, 7-[3-  
 5 [[(diphenylmethoxy)-imino]-bicyclo,[2.2.1]hept-2-yl]-; EP  
 035-rac) and 5-heptenoic acid, 7-[3-[[[(diphenylmethoxy)-  
 imino]methyl]biclo[2.2.2]-oct-5-en-2-yl]- (EP 157); and a  
 TXA<sub>2</sub> antagonist, e.g., 5-heptenoic acid, 7-[3-[[2-(phenyl-  
 amino)carbonyl]-hydrazino]methyl]7-oxabicyclo[2.2.1]hept-  
 2-yl]-, 1S[1.alpha.,2.alpha.(Z),3.alpha.,4.alpha.]]- (SQ  
 29548); benzenepropanoic acid, 2-[[3-4[(pentylamino)car-  
 10 bonyl]-2-oxazolyl]-7-oxabicyclo[2.2.1]hept-2-ylmethyl]-  
 (BMS 180291); acetic acid, [4-[2-[(phenylsulfonyl)amino]-  
 ethyl]penoxy]- (sultroban, BM-13177); benzeneacetic acid,  
 4-[2-[[[4-chlorophenyl)sulfonyl]amino]ethyl]- (daltroban,  
 BM-13505); (S-145 rac); 5-hexenoic acid, 6-[3-[[[(4-  
 15 bromophenyl)sulfonyl]amino]methyl]bicyclo[2.2.1]hep-2-  
 yl]-, decyl ester, [1S[1.alpha.2.alpha.2.alpha.(Z),-  
 3.beta.,4.alpha.]]- (ONO 8809); 9H-carbazole-9-propanoic  
 acid, 3-[[[(4-fluorophenyl)sulfonyl]amino]-1,2,3,4-tetra-  
 hydro-, (R)- (bay-u-3405); and (4Z)-6-[(5S)-5-(4-chlor-  
 20 phenylsulfonyl(aminomethyl)-cycloent-1-enyl]4-hexenoic  
 acid (ZU 154343).

Examples of combinations of active agents which can  
 be administered concurrently with a nitric oxide sub-  
 strate and/or a nitric oxide donor and a progesterone (or  
 25 other progestational agent) are low dose (e.g., 10-100 mg)  
 of aspirin (or other cyclooxygenase inhibitor; PGI<sub>2</sub>-  
 mimetics (e.g., iloprost, cicaprost); combinations of a  
 PGI<sub>2</sub>-mimetic and low dose aspirin.

Examples of dosage ranges of typical NO-substrates  
 30 and NO-donors (per os) are:

	total dose:
L-Arginine	500 mg - 10 g p.o.
Sodium Nitroprusside	range 500-2000 ug/kg/day
Nitroglycerin	0.5-10 mg
35 Isosorbid mononitrate	10-100 mg
Isosorbid dinitrate	10-100 mg

The following are typical oral dosage ranges active agents of the progestin and the optional other active agents concurrently administered with the nitric oxide substrate or donor:

- 5 Progestins: A daily dose bioequivalent to 50-300 mg of progesterone/day, e.g., an injectable suspension of medroxyprogesterone acetate to provide a weekly dose of thereof of 100-1000 mg or tablets or dragees providing an oral dose thereof of 5-10 mg/day; an  
10 injectable solution of hydroxyprogesterone caproate which provides a weekly dose of 250-500 mg; tablets, capsules or dragees of nortindrone acetate which provide a daily dose of 5-20 mg.

Cicaprost: 5-100 ug/kg/day p.o.

- 15 Aspirin: 10-100 mg/kg/day p.o.

- The pharmacologically active agents employed in this invention can be administered in admixture with conventional excipients, i.e., pharmaceutically acceptable liquid, semi-liquid or solid organic or inorganic carriers suitable, e.g., for parental or enteral application and which do not deleteriously react with the active compound in admixture therewith. Suitable pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohols, vegetable  
20 oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, pentaerythritol fatty acid esters, hydroxy methylcellulose, polyvinyl pyrrolidone, etc. The pharmaceutical  
25 preparations can be sterilized and if desired mixed with auxiliary agents, e.g., lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, coloring, flavoring and/or aromatic substances and the like which do not deleteriously react with the active compounds.  
30  
35

For parental application, particularly suitable are solutions, preferably oily or aqueous solutions, as well as suspensions, emulsions, or implants, including suppositories. Ampoules are convenient unit dosages.

In a preferred aspect, the composition of this invention is adapted for ingestion.

For enteral application, particularly suitable are unit dosage forms, e.g., tablets, dragees or capsules having talc and/or a carbohydrate carrier or binder or the like, the carrier preferably being lactose and/or corn starch and/or potato starch; particulate solids, e.g., granules; and liquids and semi-liquids, e.g., syrups and elixirs or the like, wherein a sweetened vehicle is employed. Sustained release compositions can be formulated including those wherein the active compound is protected with differentially degradable coatings, e.g., by micro-encapsulation, multiple coatings, etc.

Suitable for oral administration are, inter alia, tablets, dragees, capsules, pills, granules, suspensions and solutions. Each unit dose, e.g., each tablespoon of liquid or each tablet, or dragee contains, for example, 5-5000 mg of each active agent.

Solutions for parenteral administration contain, for example, 0.01 - 1% of each active agent in an aqueous or alcoholic solution.

The nitric oxide substrate and/or donor can be administered as an admixture with the progestational agent and any other optional active agent or as a separate unit dosage form, either simultaneously <sup>or</sup> there-with or at different times during the day from each other.

The combination of active agents is preferably administered at least once daily (unless administered in a dosage form which delivers the active agents continuously) and more preferably several times daily, e.g., in 2 to 6 divided doses. The typical dose is about 0.5 to 1000 mg of each active agent, although some less active agents, e.g., L-Arginine, require much higher oral dosages, e.g., 500 to 10,000 mg, and others, e.g., sodium nitroprusside, require lower doses, e.g., 500-2,000 ug/kg/day. Doses for nitroglycerine typically are orally 2.5 mg 2 x daily; sublingually, 0.8 mg 1-4 x daily; and



transdermally, 0.2-0.4 mg/hr. Since the LD<sub>50</sub> dosages of most of these active agents is known in the prior art, a lower dosage regimen can be initiated and the dosage increased until a positive effect is achieved or a higher dosage regimen can initially be employed, e.g., in a crisis situation, and the dosages regulated downward as relief from the symptoms is achieved.

In humans, both L-arginine and progesterone (or bioequivalent of another progestin) should be given in a ratio which produces blood plasma levels of about 1-5mMol/ml and 300-1,000 ng/ml (0.9-3μMol/l), respectively. The NO-donor, e.g., sodium nitroprusside, should be given with the progesterone (or bioequivalent of another progestin) in a ratio producing blood plasma levels of about 1-10 μMol/l and 300-1,000 ng/ml (0.9-3μMol/l), respectively.

#### Brief Description of the Drawings

With reference to the drawings,

FIGURE 1 is a series of strip chart recordings showing the effect of L-arginine on spontaneously contracting uterine strips from rat on day 18 of gestation;

FIGURE 2: Dose-dependent relaxation effects of L-arginine (0.1 mM to 10 mM) on spontaneously contracting uterine strips from rats at different stages of gestation, during delivery and post partum. The tissues were obtained on days 17-22 (d17, d18, d19 and d22) of gestation, on day 22 (d22 del) during spontaneous delivery (1-3 pups delivered), or on 1 (d1pp) and 2 (d2pp) days postpartum. The duration of complete inhibition of spontaneous uterine contraction are dose-dependent. Data are analyzed by repeated measures ANOVA on seven groups. The effects of L-arginine from concentrations of 1 mM are significantly (P<0.01) decreased during spontaneous delivery at term and postpartum, compared to all other times. Each data point represent mean ± S.E.M.

The total number of strips studied at each time period was 8-16 from 4-6 animals per group.

FIGURE 3: Dose response effects of L-arginine (0.6 mM to 10 mM) on the spontaneous contractility of uterine strips from ovariectomized adult rats. Animals received s.c. injection of 1 ug estradiol - 17b (OVX + E), 2 mg progesterone (OVX + P), estradiol and progesterone (OVX + E + P) in sesame oil or oil alone (OVX + Oil) for 3 days prior to contractility measurements. Values are mean  $\pm$  SEM for 4 strips from each animal from 4 rats per group. Data are analyzed by repeated measures ANOVA on four groups. \*P<0.05 OVX + P vs OVX + E.

FIGURE 4: 8-bromo-cGMP dose relaxation-response curves for uterine tissues from rats delivering, spontaneously at term (DEL), preterm with ZK299 (PRETERM DEL) and nondelivering (NONDEL) on day 18 of gestation. Each point represent means  $\pm$  SEM for 4 strips from each animal from 4 rats per group.

FIGURE 5 is a bar chart which shows the effect on blood pressure of test animals of 50 mg of the hypertensive agent L-NAME, alone or in combination with one or both of L-arginine and progesterone (R-5020); and

FIGURE 6 is bar chart which shows the effect in the same experiments on pup weights of these compounds.

#### 25 Discussion of the Drawings

The strip chart recordings of Figure 1 show that the application of L-arginine (1-3 mM) (A, B, E), sodium nitroprusside (5 mM) (C), nitric oxide (0.1 mM) (D) to muscle baths produced substantial relaxations. The effects of L-arginine were reversed by L-NAME (3 mM) (B) and methylene blue (0. mM) (E). These are typical recordings of 8-16 strips from 6 animals in each group. Each upstroke from baseline represents a contraction.

The strip chart recording of Figure 1C show that the application of sodium nitroprusside (SNP) caused sustained relaxation in spontaneously contracting uterine strips after a lag period and that tissues in the relax

state were responsive to potassium chloride. Similar recordings of 12 uterine strips from 4 animals were obtained.

5 The strip chart recording in Figure 1D show the relaxation produced by authentic nitric oxide gas (0.1 mM). Similar recordings were obtained from 8 strips from 4 animals.

10 The strip chart recordings of Figure 1E show that L-arginine (1 mM) produced relaxation of spontaneously contracting tissues and these effects were repeatable in the same strip (as in Fig. 1A) and that the relaxation effect of L-arginine (1 mM) was abolished by methylene blue (0.1 mM) when added before the application of L-arginine (B).

15 In the experiments whose results are shown by the graph of Figure 2, the tissues were obtained on days 17-22 (d17, d18, d19 and d22) of gestation, on day 22 (d22 del) during spontaneous delivery (1-3 pups delivered), or on 1 (d1pp) and 2 (d2pp) days postpartum. The duration of complete inhibition of spontaneous uterine contractions are dose-dependent. The effects of L-arginine from concentrations of 1 mM are significantly ( $P < 0.01$ ) decreased during spontaneous delivery at term and postpartum, compared to all other times. Each data point  
20 represent mean  $\pm$  S.E.M. The total number of strips studied at each time period was 8-16 from 4-6 animals per group.

25 In the experiments whose results are shown by the graph of Figure 3, nonpregnant ovariectomized rats received s.c. injection of 1 ug estradiol-17-B (OVX + E), 2 mg progesterone (OVX + P), estradiol and progesterone (OVX + E + P) in sesame oil or oil alone (OVX + Oil) for 3 days prior to contractility measurements. Values are mean  $\pm$  SEM for 4 strips from each animal from 4 rats per  
30 group. \* $P < 0.05$  OVX + P vs OVX + E.

35 The charge of Figure 4 shows 8-bromo-cGMP dose relaxation-response curves for uterine tissues from rats

delivering, spontaneously at term (DEL), preterm with ZK299 (PRETERM DEL) and nondelivering (NONDEL) on day 18 of gestation. Each point represent means  $\pm$  SEM for 4 strips from each animal from 4 rats per group.

5 The data in Table 1 below show the effects of L-NAME infusion on blood pressure (mm Hg) in pregnant rats.

**TABLE 1**

Gestation day	Blood Pressure (mm Hg)		
	CONTROL	L-NAME	
		25 mg/day	50 mg/day
Day 15	121 $\pm$ 3 <sup>a</sup>	119 $\pm$ 2 <sup>a</sup>	123 $\pm$ 3 <sup>a</sup>
Day 18	119 $\pm$ 3 <sup>a</sup>	144 $\pm$ 4 <sup>b</sup>	166 $\pm$ 2 <sup>c</sup>
Day 22	120 $\pm$ 5 <sup>a</sup>	146 $\pm$ 2 <sup>b</sup>	168 $\pm$ 3 <sup>c</sup>

15 Means with different superscripts differ significantly (P<0.05)

The data in Table 2 below show the delivery and the pups delivered of L-NAME infusion to pregnant rats.

**TABLE 2**

	CONTROL	L-NAME	
		25 mg/day	50 mg/day
Day of Delivery	22.3 $\pm$ 0.2	22.4 $\pm$ 0.2	22.7 $\pm$ 0.2
Total # of pups	59	65	56
# of dead pups	2	5	10
Weight of pups	6.32 $\pm$ 0.05 <sup>a</sup>	5.05 $\pm$ 0.08 <sup>b</sup>	4.56 $\pm$ 0.10 <sup>c</sup>
Total # of animals	8	9	10

25 Means with different superscripts differ significantly (P<0.05).

30 Another experiment using L-NAME-induced" pre-eclampsia" showed that treatment with L-arginine alone partially reduced blood pressure (Figure 5). Similarly, animals treated with L-NAME and R 5020 (promegestone), a progestational agent with no antimineralocorticoid effect

or other antagonistic or agonistic properties, also  
partially reduced L-NAME-induced hypertension. As also  
shown in Figure 5, when the same doses of L-arginine and  
R 5020 were given simultaneously, their combined effect  
lowered blood pressure to normal levels.

Additionally, evaluation of fetal weights in the  
same animals treated as described above, showed  
intrauterine fetal retardation (decreased weight of  
pups), typical preeclamptic fetuses (Figure 6).

Treatment of the "preeclamptic" groups of animals with  
either L-arginine alone or R 5020 alone slightly but  
statistically significant, elevated fetal weights. As  
also shown in Figure 6, the combined effect of the two  
compounds administered together significantly elevated  
fetal weight above that observed with either compound  
alone, a highly significant advantage to survival of the  
fetus under these conditions.

It can be concluded from these studies that the  
combined treatment of L-arginine with a progestational  
agent whose activity is "pure", like R 5020 provides  
results which cannot be achieved with either type of drug  
alone. The studies show that the basis for this  
effectiveness lies in the ability of the progestational  
agent to increase the effectiveness of nitric oxide (or  
L-arginine, the substrate of nitric oxide) to dilate blood  
vessels and thereby lower blood pressure as well as  
increase fetal/maternal perfusion, thereby increasing  
fetal weight.

The combined effect of the combination of these  
agents is surprisingly dramatic and, more importantly,  
the significant fetal and maternal effects observed with  
treatment with the combination. Prior medical evidence  
does not suggest that the combination would provide these  
advantages, because the basis for them is not the simple  
combination of two agonistic compounds but instead is the  
sensitizing of nitric oxide provided by the progestin.  
The studies clearly indicate that progestins increase the

effector system for nitric oxide (not increase nitric oxide synthesis).

5 The method of treatment employed in this invention can also be employed for the treatment of hypertension (in both females and males), climacteric disorders (hot flushes, mood swings) in menopausal women, thrombotic disorders, menstrual disorders (dysmenorrhea, functional uterine bleeding), and hemorrhage, etc., following the dosage regime described herein.

10 Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of  
15 the disclosure in any way whatsoever.

The entire disclosure of all applications, patents and publications, cited above and below are hereby incorporated by reference.

#### Examples

##### 20 Example 1 - Treatment of Preeclampsia

To a pregnant human female (ca 20-40 years; 60-80 kg) usually in her second half of pregnancy and displaying the symptoms of preeclampsia, including hypertension (above 140 mm systolic and above 90 mm  
25 diastolic), edema and protein-uria, administer 0.5 to 20 g of L-arginine and 200 mg of micronized progesterone per os daily in three divided doses until the symptoms are ameliorated. Thereafter, administer 0.5 to 5 mg of L-arginine and 60 mg of progesterone per os daily whenever  
30 the diastolic pressure rises above 80 mm; with increasing doses of L-arginine to from 5 to 20 mg daily until remission of the symptoms again occurs.

##### Example 2 - Treatment of preeclampsia

35 To a human female comparable to and displaying the same symptoms as the one described in Example 1, administer daily 2 x 2.5 mg of nitroglycerine and 200 mg

of progesterone following the same protocol, until the symptoms are ameliorated.

Example 3 - Treatment of Preterm Labor

5 To a human female in her sixth month of pregnancy  
and displaying symptoms of a threatened spontaneous  
abortion, including blood spotting and periodic uterine  
spasms, administer daily 17 g of L-arginine and 50 mg of  
progesterone per os daily in three divided doses until  
10 the symptoms are ameliorated. Thereafter, administer 5 g  
of L-arginine and 50 mg of progesterone per os daily with  
increasing doses to 20 g of L-arginine daily until  
remission of the symptoms again occurs.

Example 5 - Treatment of Preterm Labor

15 To a pregnant human female comparable to and  
displaying the same symptoms as the one described in  
Example 3, administer daily 2 x 25 mg of nitroglycerine  
and up to 180 mg of progesterone, following the same  
protocol, until the symptoms are ameliorated.

20 The preceding examples can be repeated with similar  
success by substituting the generically or specifically  
described reactants and/or operating conditions of this  
invention for those used in the preceding examples.

25 From the foregoing description, one skilled in the  
art can easily ascertain the essential characteristics of  
this invention, and without departing from the spirit and  
scope thereof, can make various changes and modifications  
of the invention to adapt it to various usages and  
conditions.

00121849-072428